

MATH 307: Problem Set #4

Due on: October 22, 2014

Problem 1 *Homogeneous ODEs with Const. Coeffs: Distinct Roots*

In each of the following, find the general solution of the given differential equation

(a) $y'' + 3y' + 2y = 0$

(b) $2y'' - 3y' + y = 0$

(c) $y'' - 2y' - 2y = 0$

.....

Solution 1.

- (a) The characteristic equation is $r^2 + 3r + 2$, which has roots $r_1 = -1$ and $r_2 = -2$. Hence the general solution is

$$y = C_1e^{-t} + C_2e^{-2t}.$$

- (b) The characteristic equation is $2r^2 - 3r + 1$, which has roots $r_1 = 1/2$ and $r_2 = 1$. Hence the general solution is

$$y = C_1e^{t/2} + C_2e^t.$$

- (c) The characteristic equation is $r^2 - 2r - 2$, which has roots $r_1 = 1 + \sqrt{3}$ and $r_2 = 1 - \sqrt{3}$. Hence the general solution is

$$y = C_1e^{(1+\sqrt{3})t} + C_2e^{(1-\sqrt{3})t}.$$

Problem 2 *Homogeneous IVPs with Const. Coeffs: Distinct Roots*

In each of the following, find the solution of the IVP

(a) $y'' + 4y' + 3y = 0$, $y(0) = 2$, $y'(0) = -1$

(b) $y'' + 3y' = 0$, $y(0) = -2$, $y'(0) = 3$

.....

Solution 2. (a) The characteristic equation is $r^2 + 4r + 3 = 0$ which has roots $r_1 = -1$ and $r_2 = -3$. Hence the general solution is

$$y = C_1 e^{-t} + C_2 e^{-3t}.$$

We calculate then that

$$y' = -C_1 e^{-t} - 3C_2 e^{-3t}.$$

This means that $y(0) = C_1 + C_2$ and $y'(0) = -C_1 - 3C_2$. Therefore our initial condition tells us

$$\begin{aligned} C_1 + C_2 &= 2 \\ -C_1 - 3C_2 &= -1, \end{aligned}$$

and solving this, we find $C_1 = 5/2$ and $C_2 = -1/2$. Therefore the solution is

$$y = \frac{5}{2} e^{-t} - \frac{1}{2} e^{-3t}.$$

(b) The characteristic equation is $r^2 + 3r = 0$ which has roots $r_1 = 0$ and $r_2 = -3$. Hence the general solution is

$$y = C_1 + C_2 e^{-3t}.$$

We calculate then that

$$y' = -3C_2 e^{-3t}.$$

This means that $y(0) = C_1 + C_2$ and $y'(0) = -3C_2$. Therefore our initial condition tells us

$$\begin{aligned} C_1 + C_2 &= -2 \\ -3C_2 &= 3, \end{aligned}$$

and solving this, we find $C_1 = -1$ and $C_2 = -1$. Therefore the solution is

$$y = -1 - e^{-3t}.$$

Problem 3 *Complex Number Problems*

In each of the following, rewrite the expression in the form $a + ib$

(a) e^{2-3i}

(b) $e^{2-(\pi/2)i}$

(c) π^{-1+2i}

.....

Solution 3.

(a)

$$\begin{aligned} e^{2-3i} &= e^2 e^{-3i} = e^2 (\cos(-3) + i \sin(-3)) \\ &= e^2 (\cos(3) - i \sin(3)) = e^2 \cos(3) - i e^2 \sin(3) \end{aligned}$$

(b)

$$\begin{aligned} e^{2-(\pi/2)i} &= e^2 e^{-(\pi/2)i} = e^2 (\cos(-\pi/2) + i \sin(-\pi/2)) \\ &= e^2 (0 - i) = -i e^2 \end{aligned}$$

(c)

$$\begin{aligned} \pi^{-1+2i} &= (\pi)^{-1+2i} = (e^{\ln(\pi)})^{-1+2i} = e^{\ln(\pi)(-1+2i)} \\ &= e^{-\ln(\pi)+2\ln(\pi)i} = e^{-\ln(\pi)} e^{2\ln(\pi)i} \\ &= e^{-\ln(\pi)} (\cos(2 \ln(\pi)) + i \sin(2 \ln(\pi))) \\ &= e^{-\ln(\pi)} \cos(2 \ln(\pi)) + i e^{-\ln(\pi)} \sin(2 \ln(\pi)) \\ &= \frac{1}{\pi} \cos(2 \ln(\pi)) + i \frac{1}{\pi} \sin(2 \ln(\pi)) \end{aligned}$$

Problem 4 *Homogeneous ODEs with Const. Coeffs: Complex Roots*

In each of the following, find the general solution of the ODE

(a) $y'' - 2y' + 6y = 0$

(b) $y'' + 2y' + 2y = 0$

(c) $y'' + 4y' + 6.25y = 0$

.....

Solution 4.

- (a) The corresponding characteristic equation is $r^2 - 2r + 6 = 0$, which has roots $1 \pm \sqrt{5}i$. Hence the general solutions is

$$y = C_1 e^t \cos(\sqrt{5}t) + C_2 e^t \sin(\sqrt{5}t).$$

- (b) The corresponding characteristic equation is $r^2 + 2r + 2 = 0$, which has roots $r_1 = -1 \pm i$. Hence the general solutions is

$$y = C_1 e^{-t} \cos(t) + C_2 e^{-t} \sin(t).$$

- (c) The corresponding characteristic equation is $r^2 + 4r + 6.25 = 0$, which has roots $r_1 = -2 \pm \frac{3}{2}i$. Hence the general solutions is

$$y = C_1 e^{-2t} \cos(3t/2) + C_2 e^{-2t} \sin(3t/2).$$

Problem 5 *Homogeneous IVPs with Const. Coeffs: Complex Roots*

In each of the following, find the solution of the IVP

- (a) $y'' + 4y = 0, \quad y(0) = 0, \quad y'(0) = 1$
 (b) $y'' + 4y' + 5y = 0, \quad y(0) = 1, \quad y'(0) = 0$

.....

Solution 5.

- (a) The corresponding characteristic equation is $r^2 + 4 = 0$, which has roots $\pm 2i$. Hence the general solution is

$$y = C_1 \cos(2t) + C_2 \sin(2t).$$

Therefore

$$y' = -2C_1 \sin(2t) + 2C_2 \cos(2t),$$

and it follows that $y(0) = C_1$ and $y'(0) = 2C_2$. Then our initial condition tells us

$$\begin{aligned} C_1 &= 0 \\ 2C_2 &= 1 \end{aligned}$$

and therefore $C_1 = 0$ and $C_2 = 1/2$, so that the solution to the initial value problem is

$$y = \frac{1}{2} \sin(2t).$$

- (b) The corresponding characteristic equation is $r^2 + 4r + 5 = 0$, which has roots $-2 \pm i$. Hence the general solution is

$$y = C_1 e^{-2t} \cos(t) + C_2 e^{-2t} \sin(t).$$

Therefore

$$y' = -2C_1 e^{-2t} \cos(t) - C_1 e^{-2t} \sin(t) - 2C_2 e^{-2t} \sin(t) + C_2 e^{-2t} \cos(t),$$

and it follows that $y(0) = C_1$ and $y'(0) = -2C_1 + C_2$. Then our initial condition tells us

$$\begin{aligned} C_1 &= 1 \\ -2C_1 + C_2 &= 0 \end{aligned}$$

and therefore $C_1 = 1$ and $C_2 = 2$, so that the solution to the initial value problem is

$$y = e^{-2t} \cos(t) + 2e^{-2t} \sin(t).$$

Problem 6 *Homogeneous ODEs with Const. Coeffs: Repeated Roots*

In each of the following, find the general solution of the ODE

- (a) $9y'' + 6y' + y = 0$
- (b) $4y'' + 12y' + 9y = 0$
- (c) $y'' - 6y' + 9y = 0$
- (d) $25y'' - 20y' + 4y = 0$

.....

Solution 6.

- (a) The roots of the characteristic equation are $r_1 = r_2 = -1/3$, and therefore the general solution is

$$y = C_1 e^{-t/3} + C_2 t e^{-t/3}.$$

- (b) The roots of the characteristic equation are $r_1 = r_2 = -3/2$, and therefore the general solution is

$$y = C_1 e^{-3t/2} + C_2 t e^{-3t/2}.$$

- (c) The roots of the characteristic equation are $r_1 = r_2 = 3$, and therefore the general solution is

$$y = C_1 e^{3t} + C_2 t e^{3t}.$$

- (d) The roots of the characteristic equation are $r_1 = r_2 = 2/5$, and therefore the general solution is

$$y = C_1 e^{2t/5} + C_2 t e^{2t/5}.$$

Problem 7 *Reduction of Order*

In each of the following, use the method of reduction of order to find a second solution of the ODE

- (a) $t^2y'' + 2ty' - 2y = 0$, $t > 0$ (one solution is $y(t) = t$)
 (b) $(x - 1)y'' - xy' + y = 0$, $x > 1$ (one solution is $y(x) = e^x$)

.....

Solution 7.

- (a) We try a solution of the form $y = v(t)t$. Then $y' = v'(t)t + v(t)$ and $y''(t) = v''(t)t + 2v'(t)$, so that

$$\begin{aligned} t^2y'' + 2ty' - 2y &= t^2(v''(t)t + 2v'(t)) + 2t(v'(t)t + v(t)) - 2(v(t)t) \\ &= t^3v''(t) + 4t^2v'(t). \end{aligned}$$

Then since $t^2y'' + 2ty' - 2y = 0$ (in order to be a solution to the equation), we must have

$$t^3v''(t) + 4t^2v'(t) = 0.$$

Dividing both sides by t^2 , this simplifies to

$$tv''(t) + 4v'(t) = 0.$$

Now if we substitute $w = v'$, then this equation becomes

$$tw'(t) + 4w(t) = 0.$$

This equation is separable, and the solution is $w(t) = C_1t^{-4}$, where C_1 is an arbitrary constant. Then since $v'(t) = w$, it follows that $v(t) = C_1t^{-3} + C_2$ (where we've left $-C_1/3$ as C_1 since it's an arbitrary constant anyway). Hence another solution is

$$y = v(t)t = C_1t^{-2} + C_2t,$$

and in fact this is the general solution.

- (b) We try a solution of the form $y = v(x)e^x$. Then $y' = v'(x)e^x + v(x)e^x$ and $y'' = v''(x)e^x + 2v'(x)e^x + v(x)e^x$, so that

$$\begin{aligned} (x - 1)y'' - xy' + y &= (x - 1)(v''(x)e^x + 2v'(x)e^x + v(x)e^x) - x(v'(x)e^x + v(x)e^x) + (v(x)e^x) \\ &= (x - 1)e^xv''(x) + (x - 2)e^xv'(x). \end{aligned}$$

Then since $(x - 1)y'' - xy' + y = 0$ (in order to be a solution to the equation), we must have

$$(x - 1)e^xv''(x) + (x - 2)e^xv'(x) = 0.$$

Dividing both sides by e^x , this simplifies to

$$(x - 1)v''(x) + (x - 2)v'(x) = 0.$$

Now if we substitute $w = v'$, the equation becomes

$$(x - 1)w'(x) + (x - 2)w(x) = 0,$$

which is separable. The solution is

$$w = C_1(x - 1)e^{-x}.$$

Then since $v' = w$, it follows that

$$v = -C_1xe^{-x} + C_2.$$

Hence another solution to the original differential equation is

$$y = ve^x = -C_1x + C_2e^x,$$

and in fact this is the general solution.