

MATH 307: Problem Set #7

Due on: November 3, 2014

Problem 1 *Inverse Laplace Transforms*

In each of the following, find the inverse Laplace transform of the given function

(a) $F(s) = \frac{3}{s^2+4}$

(b) $F(s) = \frac{4}{(s-1)^3}$

(c) $F(s) = \frac{2}{s^2+3s-4}$

(d) $F(s) = \frac{3s}{s^2-s-6}$

(e) $F(s) = \frac{2s+1}{s^2-2s+2}$

(f) $F(s) = \frac{8s^2-4s+12}{s(s^2+4)}$

(g) $F(s) = \frac{1-2s}{s^2+4s+5}$

(h) $F(s) = \frac{2s-3}{s^2+2s+10}$

.....

Problem 2 *Solving IVPs*

In each of the following, use Laplace transforms to solve the initial value problem

(a) $y'' + 3y' + 2y = 0, y(0) = 1, y'(0) = 0$

(b) $y'' - 2y' + 2y = 0, y(0) = 0, y'(0) = 1$

(c) $y'' + 2y' + 5y = 0, y(0) = 2, y'(0) = -1$

(d) $y'' + \omega^2 y = \cos(2t), y(0) = 1, y'(0) = 0$ (Here, ω is a constant and $\omega^2 \neq 4$.)

.....

Problem 3 *Step Functions*

In each of the following, sketch a graph of the given function on the interval $t \geq 0$

(a) $g(t) = u_1(t) + 2u_3(t) - 6u_4(t)$

(b) $g(t) = (t - 1)u_1(t) - 2(t - 2)u_2(t) + (t - 3)u_3(t)$

(c)

.....

Problem 4 *Brackets to Step Functions*

In each of the following, sketch a graph of the given function on the interval $t \geq 0$, and then convert the function from bracket form to step function form

(a)

$$f(t) = \begin{cases} 1 & \text{if } 0 \leq t < 2 \\ e^{-(t-2)} & \text{if } t \geq 2 \end{cases}$$

(b)

$$f(t) = \begin{cases} t & \text{if } 0 \leq t < 1 \\ t - 1 & \text{if } 1 \leq t < 2 \\ t - 2 & \text{if } 2 \leq t < 3 \\ 0 & \text{if } t \geq 3 \end{cases}$$

.....

Problem 5 *Laplace Transforms of Step Functions*

Find the Laplace transform of the given function

(a)

$$f(t) = \begin{cases} 0 & \text{if } 0 \leq t < 2 \\ (t - 2)^2 & \text{if } t \geq 2 \end{cases}$$

(b)

$$f(t) = \begin{cases} 0 & \text{if } 0 \leq t < 1 \\ t^2 - 2t + 2 & \text{if } t \geq 1 \end{cases}$$

.....

Problem 6 *More Inverse Laplace Transforms*

Find the Inverse Laplace transform of the given function

$$(a) F(s) = \frac{e^{-2s}}{s^2+s-2}$$

$$(b) F(s) = \frac{2e^{-2s}}{s^2-4}$$

$$(c) F(s) = \frac{e^{-s}+e^{-2s}-e^{-3s}-e^{-4s}}{s}$$

.....

Problem 7 *Solutions to IVPs with Discontinuous Forcing*

In each of the following problems, find the solution of the given initial value problem

$$(a) y'' + 2y' + 2y = h(t), y(0) = 0, y'(0) = 1, h(t) = \begin{cases} 0 & \text{if } 0 \leq t < \pi \\ 1 & \text{if } \pi \leq t < 2\pi \\ 0 & \text{if } 2\pi \leq t \end{cases}$$

$$(b) y'' + 3y' + 2y = f(t), y(0) = 0, y'(0) = 0, f(t) = \begin{cases} 1 & \text{if } 0 \leq t < 10 \\ 0 & \text{if } t \geq 10 \end{cases}$$

$$(c) y'' + y = u_{3\pi}(t), y(0) = 1, y'(0) = 0.$$

.....