

Math 307 Lecture 9

Exact Equations and Integrating Factors Part II

W.R. Casper

Department of Mathematics
University of Washington

April 18, 2014

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Today!

Last time:

- Numerical Approximations

This time:

- Return of Exact Equations and Integrating Factors!

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Outline

- 1 Exact Equations
 - Review of Basic Definitions
 - Why do we Like Exact Equations?
 - Solving Exact Equations
- 2 Integrating Factors
 - Integrating Factor Review
 - Integrating Factor Examples

Outline

- 1 **Exact Equations**
 - Review of Basic Definitions
 - Why do we Like Exact Equations?
 - Solving Exact Equations
- 2 **Integrating Factors**
 - Integrating Factor Review
 - Integrating Factor Examples

What is an Exact Equation?

- Remember that any first-order ODE can be written in the form

$$M(x, y) + N(x, y)y' = 0.$$

- Recall the definition of an exact equation:

Definition

An equation of the above form is *exact* if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- That's well and good, but let's see some examples!

What is an Exact Equation?

- Remember that any first-order ODE can be written in the form

$$M(x, y) + N(x, y)y' = 0.$$

- Recall the definition of an exact equation:

Definition

An equation of the above form is *exact* if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- That's well and good, but let's see some examples!

What is an Exact Equation?

- Remember that any first-order ODE can be written in the form

$$M(x, y) + N(x, y)y' = 0.$$

- Recall the definition of an exact equation:

Definition

An equation of the above form is *exact* if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- That's well and good, but let's see some examples!

What is an Exact Equation?

- Remember that any first-order ODE can be written in the form

$$M(x, y) + N(x, y)y' = 0.$$

- Recall the definition of an exact equation:

Definition

An equation of the above form is *exact* if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- That's well and good, but let's see some examples!

What is an Exact Equation?

- Remember that any first-order ODE can be written in the form

$$M(x, y) + N(x, y)y' = 0.$$

- Recall the definition of an exact equation:

Definition

An equation of the above form is *exact* if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- That's well and good, but let's see some examples!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppies!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppies!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppies!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppies!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppers!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppers!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppers!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppers!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

What is an Exact Equation?

In each of the following, let's try to decide if the equation is exact

- $y \cos(xy) + x \cos(xy)y' = 0$
- Yes!
- $y \cos(xy) - \tan(x) + x \cos(xy)y' = 0$
- Yuppers!
- $2xy + x^2yy' = 0$
- No sir!
- $ye^{xy} + (xe^{xy} + \sec^2(y))y' = 0$
- You 'betcha!

Outline

- 1 **Exact Equations**
 - Review of Basic Definitions
 - **Why do we Like Exact Equations?**
 - Solving Exact Equations

- 2 **Integrating Factors**
 - Integrating Factor Review
 - Integrating Factor Examples

Why are Exact Equations Nice?

- The next theorem tells us why we **love** exact equations:

Theorem

Suppose that the equation

$$M(x, y) + N(x, y)y' = 0$$

is exact and that M, N are “nice enough”. Then there exists a function $\psi(x, y)$ such that

$$\frac{\partial \psi}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial \psi}{\partial y} = N(x, y)$$

Why are Exact Equations Nice?

- The next theorem tells us why we **love** exact equations:

Theorem

Suppose that the equation

$$M(x, y) + N(x, y)y' = 0$$

is exact and that M, N are “nice enough”. Then there exists a function $\psi(x, y)$ such that

$$\frac{\partial \psi}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial \psi}{\partial y} = N(x, y)$$

Why are Exact Equations Nice?

- The next theorem tells us why we **love** exact equations:

Theorem

Suppose that the equation

$$M(x, y) + N(x, y)y' = 0$$

is exact and that M, N are “nice enough”. Then there exists a function $\psi(x, y)$ such that

$$\frac{\partial \psi}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial \psi}{\partial y} = N(x, y)$$

Why are Exact Equations Nice?

- It may not look like it yet, but the previous theorem helps us to solve exact equations!
- To see why, let's calculate $\frac{d\psi}{dx}$ using implicit differentiation:

$$\begin{aligned}\frac{d\psi(x, y)}{dx} &= \frac{\partial\psi}{\partial x} \frac{dx}{dx} + \frac{\partial\psi}{\partial y} \frac{dy}{dx} \\ &= \frac{\partial\psi}{\partial x} + \frac{\partial\psi}{\partial y} y' \\ &= M(x, y) + N(x, y)y'\end{aligned}$$

Why are Exact Equations Nice?

- It may not look like it yet, but the previous theorem helps us to solve exact equations!
- To see why, let's calculate $\frac{d\psi}{dx}$ using implicit differentiation:

$$\begin{aligned}\frac{d\psi(x, y)}{dx} &= \frac{\partial\psi}{\partial x} \frac{dx}{dx} + \frac{\partial\psi}{\partial y} \frac{dy}{dx} \\ &= \frac{\partial\psi}{\partial x} + \frac{\partial\psi}{\partial y} y' \\ &= M(x, y) + N(x, y)y'\end{aligned}$$

Why are Exact Equations Nice?

- It may not look like it yet, but the previous theorem helps us to solve exact equations!
- To see why, let's calculate $\frac{d\psi}{dx}$ using implicit differentiation:

$$\begin{aligned}\frac{d\psi(x, y)}{dx} &= \frac{\partial\psi}{\partial x} \frac{dx}{dx} + \frac{\partial\psi}{\partial y} \frac{dy}{dx} \\ &= \frac{\partial\psi}{\partial x} + \frac{\partial\psi}{\partial y} y' \\ &= M(x, y) + N(x, y)y'\end{aligned}$$

Why are Exact Equations Nice?

- It may not look like it yet, but the previous theorem helps us to solve exact equations!
- To see why, let's calculate $\frac{d\psi}{dx}$ using implicit differentiation:

$$\begin{aligned}\frac{d\psi(x, y)}{dx} &= \frac{\partial\psi}{\partial x} \frac{dx}{dx} + \frac{\partial\psi}{\partial y} \frac{dy}{dx} \\ &= \frac{\partial\psi}{\partial x} + \frac{\partial\psi}{\partial y} y' \\ &= M(x, y) + N(x, y)y'\end{aligned}$$

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Why are Exact Equations Nice?

- Thus the differential equation

$$M(x, y) + N(x, y)y' = 0$$

becomes the equation

$$\frac{d\psi(x, y)}{dx} = 0$$

- Integrating both sides with respect to x , the solution is then

$$\psi(x, y) = C$$

- We've solved the exact equation!!

Outline

- 1 **Exact Equations**
 - Review of Basic Definitions
 - Why do we Like Exact Equations?
 - **Solving Exact Equations**

- 2 **Integrating Factors**
 - Integrating Factor Review
 - Integrating Factor Examples

Solving Exact Equations

- We've found that the solution to an exact equation is given implicitly by $\psi(x, y) = C$
- One important question you should be asking yourself now is how do we determine $\psi(x, y)$?
- The answer is "partial integration"
- To see what we mean, let's look at some examples!

Solving Exact Equations

- We've found that the solution to an exact equation is given implicitly by $\psi(x, y) = C$
- One important question you should be asking yourself now is how do we determine $\psi(x, y)$?
- The answer is "partial integration"
- To see what we mean, let's look at some examples!

Solving Exact Equations

- We've found that the solution to an exact equation is given implicitly by $\psi(x, y) = C$
- One important question you should be asking yourself now is how do we determine $\psi(x, y)$?
- The answer is "partial integration"
- To see what we mean, let's look at some examples!

Solving Exact Equations

- We've found that the solution to an exact equation is given implicitly by $\psi(x, y) = C$
- One important question you should be asking yourself now is how do we determine $\psi(x, y)$?
- The answer is "partial integration"
- To see what we mean, let's look at some examples!

Solving Exact Equations

- We've found that the solution to an exact equation is given implicitly by $\psi(x, y) = C$
- One important question you should be asking yourself now is how do we determine $\psi(x, y)$?
- The answer is "partial integration"
- To see what we mean, let's look at some examples!

Solving Exact Equations: A First Example

- Consider the equation

$$\overbrace{y \cos(xy) - \tan(x)}^{M(x,y)} + \overbrace{x \cos(xy)}^{N(x,y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial y} = N(x, y) = x \cos(xy).$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial y} \partial y = \int x \cos(xy) \partial y$$

Solving Exact Equations: A First Example

- Consider the equation

$$\overbrace{y \cos(xy) - \tan(x)}^{M(x,y)} + \overbrace{x \cos(xy)}^{N(x,y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial y} = N(x, y) = x \cos(xy).$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial y} \partial y = \int x \cos(xy) \partial y$$

Solving Exact Equations: A First Example

- Consider the equation

$$\overbrace{y \cos(xy) - \tan(x)}^{M(x,y)} + \overbrace{x \cos(xy)}^{N(x,y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial y} = N(x, y) = x \cos(xy).$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial y} \partial y = \int x \cos(xy) \partial y$$

Solving Exact Equations: A First Example

- Consider the equation

$$\overbrace{y \cos(xy) - \tan(x)}^{M(x,y)} + \overbrace{x \cos(xy)}^{N(x,y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial y} = N(x, y) = x \cos(xy).$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial y} \partial y = \int x \cos(xy) \partial y$$

Solving Exact Equations: A First Example

- Consider the equation

$$\overbrace{y \cos(xy) - \tan(x)}^{M(x,y)} + \overbrace{x \cos(xy)}^{N(x,y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial y} = N(x, y) = x \cos(xy).$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial y} \partial y = \int x \cos(xy) \partial y$$

Solving Exact Equations: A First Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int x \cos(xy) dy$$

- To do the partial integral wrt. y , you treat x as a constant:

$$\psi(x, y) = \overbrace{\sin(xy)}^{\text{integral with } x \text{ constant}} + \overbrace{g(x)}^{\text{arbit. func. of integ.}}$$

- With partial integrals, we end up with an arbitrary “function of integration” instead of an arbitrary constant
- Notice we integrated wrt. y so we get an arbitrary func of x

Solving Exact Equations: A First Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int x \cos(xy) \partial y$$

- To do the partial integral wrt. y , you treat x as a constant:

$$\psi(x, y) = \overbrace{\sin(xy)}^{\text{integral with } x \text{ constant}} + \overbrace{g(x)}^{\text{arbit. func. of integ.}}$$

- With partial integrals, we end up with an arbitrary "function of integration" instead of an arbitrary constant
- Notice we integrated wrt. y so we get an arbitrary func of x

Solving Exact Equations: A First Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int x \cos(xy) \partial y$$

- To do the partial integral wrt. y , you treat x as a constant:

$$\psi(x, y) = \overbrace{\sin(xy)}^{\text{integral with } x \text{ constant}} + \overbrace{g(x)}^{\text{arbit. func. of integ.}}$$

- With partial integrals, we end up with an arbitrary “function of integration” instead of an arbitrary constant
- Notice we integrated wrt. y so we get an arbitrary func of x

Solving Exact Equations: A First Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int x \cos(xy) \partial y$$

- To do the partial integral wrt. y , you treat x as a constant:

$$\psi(x, y) = \overbrace{\sin(xy)}^{\text{integral with } x \text{ constant}} + \overbrace{g(x)}^{\text{arbit. func. of integ.}}$$

- With partial integrals, we end up with an arbitrary “function of integration” instead of an arbitrary constant
- Notice we integrated wrt. y so we get an arbitrary func of x

Solving Exact Equations: A First Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int x \cos(xy) \partial y$$

- To do the partial integral wrt. y , you treat x as a constant:

$$\psi(x, y) = \overbrace{\sin(xy)}^{\text{integral with } x \text{ constant}} + \overbrace{g(x)}^{\text{arbit. func. of integ.}}$$

- With partial integrals, we end up with an arbitrary “function of integration” instead of an arbitrary constant
- Notice we integrated wrt. y so we get an arbitrary func of x

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that
 $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that
 $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that
 $g(x) = \ln |\cos(x)|$ (we can forget the constant)

- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A First Example

- How can we figure out what $g(x)$ must be?
- Remember that

$$\partial\psi/\partial x = M(x, y)$$

- Therefore

$$y \cos(xy) + g'(x) = y \cos(xy) - \tan(x).$$

- This simplifies to $g'(x) = -\tan(x)$, so that $g(x) = \ln |\cos(x)|$ (we can forget the constant)
- Thus

$$\psi(x, y) = \sin(xy) + \ln |\cos(x)|$$

- Solution is therefore $\sin(xy) + \ln |\cos(x)| = C$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\overbrace{ye^{xy}}^{M(x,y)} + \overbrace{xe^{xy} + \sec^2(y)y'}^{N(x,y)} = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\overbrace{ye^{xy}}^{M(x,y)} + \overbrace{xe^{xy} + \sec^2(y)y'}^{N(x,y)} = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\underbrace{M(x,y)}_{ye^{xy}} + \underbrace{N(x,y)}_{xe^{xy} + \sec^2(y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\underbrace{M(x,y)}_{ye^{xy}} + \underbrace{N(x,y)}_{xe^{xy} + \sec^2(y)} y' = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\overbrace{ye^{xy}}^{M(x,y)} + \overbrace{xe^{xy} + \sec^2(y)y'}^{N(x,y)} = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Let's look at another example!
- Consider the equation

$$\overbrace{ye^{xy}}^{M(x,y)} + \overbrace{xe^{xy} + \sec^2(y)y'}^{N(x,y)} = 0$$

- This equation is exact (check this!)
- This means that there is a $\psi(x, y)$ satisfying

$$\frac{\partial \psi(x, y)}{\partial x} = M(x, y) = ye^{xy}.$$

- Doing a partial integral of both sides,

$$\int \frac{\partial \psi(x, y)}{\partial x} \partial x = \int ye^{xy} \partial x$$

Solving Exact Equations: A Second Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int ye^{xy} \partial x$$

- To do the partial integral wrt. x , you treat y as a constant:

$$\psi(x, y) = e^{xy} + h(y)$$

- With partial integrals, we end up with an arbitrary “function of integration” $h(y)$ instead of an arbitrary constant
- Notice we integrated wrt. x so we get an arbitrary func of y

Solving Exact Equations: A Second Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int ye^{xy} \partial x$$

- To do the partial integral wrt. x , you treat y as a constant:

$$\psi(x, y) = e^{xy} + h(y)$$

- With partial integrals, we end up with an arbitrary “function of integration” $h(y)$ instead of an arbitrary constant
- Notice we integrated wrt. x so we get an arbitrary func of y

Solving Exact Equations: A Second Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int ye^{xy} \partial x$$

- To do the partial integral wrt. x , you treat y as a constant:

$$\psi(x, y) = e^{xy} + h(y)$$

- With partial integrals, we end up with an arbitrary “function of integration” $h(y)$ instead of an arbitrary constant
- Notice we integrated wrt. x so we get an arbitrary func of y

Solving Exact Equations: A Second Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int ye^{xy} \partial x$$

- To do the partial integral wrt. x , you treat y as a constant:

$$\psi(x, y) = e^{xy} + h(y)$$

- With partial integrals, we end up with an arbitrary “function of integration” $h(y)$ instead of an arbitrary constant
- Notice we integrated wrt. x so we get an arbitrary func of y

Solving Exact Equations: A Second Example

- Doing a partial integral of both sides,

$$\psi(x, y) = \int ye^{xy} \partial x$$

- To do the partial integral wrt. x , you treat y as a constant:

$$\psi(x, y) = e^{xy} + h(y)$$

- With partial integrals, we end up with an arbitrary “function of integration” $h(y)$ instead of an arbitrary constant
- Notice we integrated wrt. x so we get an arbitrary func of y

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$ (we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$ (we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$
(we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$
(we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$ (we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$ (we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Solving Exact Equations: A Second Example

- How can we figure out what $h(y)$ must be?
- Remember that

$$\partial\psi/\partial y = N(x, y)$$

- Therefore

$$xe^{xy} + h'(y) = xe^{xy} + \sec^2(y).$$

- This simplifies to $h'(y) = \sec^2(y)$, so that $h(y) = \tan(y)$ (we can forget the constant)
- Thus

$$\psi(x, y) = e^{xy} + \tan(y)$$

- Solution is therefore $e^{xy} + \tan(y) = C$

Outline

- 1 Exact Equations
 - Review of Basic Definitions
 - Why do we Like Exact Equations?
 - Solving Exact Equations
- 2 Integrating Factors
 - Integrating Factor Review
 - Integrating Factor Examples

What if the Equation is not Exact?

Figure : A graphical depiction of the ecstasy one feels when the nonlinear first order equation is exact



What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible* in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible** in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible* in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible* in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible* in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

Figure : Eeyore never gets exact equations on exams



- "most" nonlinear first order equations are not exact
- Q: what should we do with such an equation?
- A: try to find an integrating factor!
- this will be *totally impossible* in general
- but it will work often enough to make it worth a try...

What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
 - eg. $\mu(x, y) = \mu(x)$
 - or $\mu(x, y) = \mu(y)$
 - or $\mu(x, y) = x^a y^b$
 - these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



What if the Equation is not Exact?

- Remember, an integrating factor is a function $\mu(x, y)$ that we multiply by to make the equation exact
- When we try to find one, we often make an assumption about the form
- eg. $\mu(x, y) = \mu(x)$
- or $\mu(x, y) = \mu(y)$
- or $\mu(x, y) = x^a y^b$
- these might not work; μ may be too hard to find!

Figure : Eeyore getting help from his turtle friend to find an integrating factor



Outline

- 1 Exact Equations
 - Review of Basic Definitions
 - Why do we Like Exact Equations?
 - Solving Exact Equations

- 2 Integrating Factors
 - Integrating Factor Review
 - Integrating Factor Examples

Integrating Factor Example 1

Example

Solve the first order equation

$$y + (2x - ye^y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(y)$.

- $\mu(y)y + (2x - ye^y)\mu(y)y' = 0$ exact
- Implies $\mu'(y)y + \mu(y) = 2\mu(y)$
- Results in $\mu'(y)y = \mu(y)$; a solution is $\mu(y) = y$

Integrating Factor Example 1

Example

Solve the first order equation

$$y + (2x - ye^y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(y)$.

- $\mu(y)y + (2x - ye^y)\mu(y)y' = 0$ exact
- Implies $\mu'(y)y + \mu(y) = 2\mu(y)$
- Results in $\mu'(y)y = \mu(y)$; a solution is $\mu(y) = y$

Integrating Factor Example 1

Example

Solve the first order equation

$$y + (2x - ye^y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(y)$.

- $\mu(y)y + (2x - ye^y)\mu(y)y' = 0$ exact
- Implies $\mu'(y)y + \mu(y) = 2\mu(y)$
- Results in $\mu'(y)y = \mu(y)$; a solution is $\mu(y) = y$

Integrating Factor Example 1

Example

Solve the first order equation

$$y + (2x - ye^y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(y)$.

- $\mu(y)y + (2x - ye^y)\mu(y)y' = 0$ exact
- Implies $\mu'(y)y + \mu(y) = 2\mu(y)$
- Results in $\mu'(y)y = \mu(y)$; a solution is $\mu(y) = y$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
 $2xy + h'(y) = 2xy - y^2 e^y$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 1

- Thus we get an exact equation

$$y^2 + (2xy - y^2 e^y)y' = 0$$

- We know $\psi(x, y) = \int y^2 \partial x = xy^2 + h(y)$
- Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we also know
$$2xy + h'(y) = 2xy - y^2 e^y$$
- Hence $h = \int -y^2 e^y dy = -(y^2 - 2y + 2)e^y$
- Thus $\psi(x, y) = xy^2 - (y^2 - 2y + 2)e^y$
- Solution is $\psi(x, y) = C$, ie.

$$xy^2 - (y^2 - 2y + 2)e^y = C$$

Integrating Factor Example 2

Example

Solve the first order equation

$$(x + 2) \sin(y) + x \cos(y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(x)$.

- Try it yourself!

Integrating Factor Example 2

Example

Solve the first order equation

$$(x + 2) \sin(y) + x \cos(y)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = \mu(x)$.

- Try it yourself!

Integrating Factor Example 3

Example

Solve the first order equation

$$x^2 y^3 + x(1 + y^2)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = x^a y^b$ for some constants a, b

- Try it yourself!

Integrating Factor Example 3

Example

Solve the first order equation

$$x^2y^3 + x(1 + y^2)y' = 0$$

by finding an integrating factor of the form $\mu(x, y) = x^a y^b$ for some constants a, b

- Try it yourself!

Review!

Today:

- More on Exact Equations and Integrating Factors

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Review!

Today:

- More on Exact Equations and Integrating Factors

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Review!

Today:

- More on Exact Equations and Integrating Factors

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients

Review!

Today:

- More on Exact Equations and Integrating Factors

Next time:

- Higher-order Homogeneous Linear Equations with Constant Coefficients