MATH 309: Homework #1

Due on: April 8, 2016

Problem 1 Matrix Algebra

Let A and B be the matrices

$$A = \begin{pmatrix} 1 & -1 & 1 \\ -1 & 1 & 1 \\ 1 & 1 & -1 \end{pmatrix} \quad B = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

Determine values of the following

- $\bullet AB$
- $\bullet BA$
- 3A + 2B

• • • • • • • • •

Problem 2 Matrix Puzzles

- (a) Find a nonzero square matrix A with $A^2 = 0$ (here 0 means the zero matrix)
- (b) Find a square matrix J with real entries satisfying $J^2 = -I$
- (c) Find an invertible matrix P with $P^{-1} = P^{\dagger}$ (here P^{\dagger} refers to the conjugate transpose of P)
- (d) Find a nonzero square matrix P with $P^2 = P$
- (e) Find all possible 2×2 matrices X satisfying the equation $X^2 2X + I = 0$

• • • • • • • • •

Problem 3 Matrix Inverses

For each of the following matrices, find the inverse of the matrix or explain why it doesn't have one

(a)
$$\begin{pmatrix} 1 & 2 \\ 3 & 1 \\ 4 & 7 \end{pmatrix}$$

(b) $\begin{pmatrix} 2 & -4 \\ 3 & -1 \end{pmatrix}$
(c) $\begin{pmatrix} 1 & 2 & 7 \\ 3 & 4 & 1 \\ 7 & 10 & 9 \end{pmatrix}$
(d) $\begin{pmatrix} 1 & 3 & 0 \\ 3 & 2 & 1 \\ 5 & 2 & 5 \end{pmatrix}$

Problem 4 Linear Systems

For each of the following linear systems of equations, either find the general solution, or show that no solution exists.

.

(a)

$$2x_1 + 3x_2 + x_3 = 1$$
$$x_1 + x_2 - 2x_3 = 2$$

(b)

$$x_1 + 3x_2 = 0$$

$$3x_1 + 2x_2 + x_3 = 1$$

$$5x_1 + 2x_2 + 5x_3 = 1$$

(c)

$$x_1 + 2x_2 + 7x_3 = 1$$

$$3x_1 + 4x_2 + x_3 = 0$$

$$7x_1 + 10x_2 + 9x_3 = 1$$

(d)

$$x_1 + 2x_2 + 7x_3 = 0$$

$$3x_1 + 4x_2 + x_3 = 0$$

$$7x_1 + 10x_2 + 9x_3 = 0$$

.

Problem 5 Eigenvectors and Eigenvalues

For each of the following matrices, determine the following information

- (i) the eigenvalues
- (ii) the algebraic and geometric multiplicity of each eigenvalue
- (iii) a basis for the eigenspace of each eigenvalue

(a) $\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ (b) $\begin{pmatrix} 0 & 1 \\ 1 & -1 \end{pmatrix}$ (c) $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ (d) $\begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & -2 \\ 3 & 2 & 1 \end{pmatrix}$ (e) $\begin{pmatrix} 3 & 2 & 2 \\ 1 & 4 & 1 \\ -2 & -4 & -1 \end{pmatrix}$

Problem 6 Eigenvectors and Linear Independence

Suppose that A is an $n \times n$ matrix and that \vec{v}_1 and \vec{v}_2 are eigenvectors of A with eigenvalues λ_1 and λ_2 , respectively. Show that if $\lambda_1 \neq \lambda_2$, then \vec{v}_1 and \vec{v}_2 must be linearly independent. (Here, by "show", we mean make a formal argument using both math and complete sentences.)

.

.

• • • • • • • • •

Problem 7 First Order Homogeneous Linear Systems of Ordinary Differential Equations with Constant Coefficients

Find the general solution of each of the following systems of first order homogeneous linear ordinary differential equations with constant coefficients

(a)

$$\begin{aligned}
 x_1' &= x_1 + x_2 \\
 x_2' &= x_1 + 2x_2
 \end{aligned}$$

(b)

(c)

$$x'_{1} = 3x_{1} + x_{2}$$
$$x'_{2} = 2x_{1} + 2x_{2}$$
$$x'_{1} = x_{1} + x_{2}$$
$$x'_{2} = x_{2}$$
....

Problem 8 Solution Space

Let $A(x) = (a_{ij}(x))$ be an $n \times n$ matrix, with the functions $a_{ij}(x)$ continuous on the interval (α, β) for all i, j. Consider the differential equation

$$\vec{y}'(x) = A(x)\vec{y}(x).$$

- (a) Explain why the set of solutions to this equation on the interval (α, β) is a vector space
- (b) Explain why the dimension of the solution space on the interval (α, β) is *n*-dimensional (I am asking you to reproduce the argument we did in lecture)

• • • • • • • • •

Problem 9 Wronskian Issues

Let $A(x) = (a_{ij}(x))$ be an $n \times n$ matrix, with the functions $a_{ij}(x)$ continuous on the interval (α, β) for all i, j. Consider the differential equation

$$\vec{y}'(x) = A(x)\vec{y}(x).$$

Recall that the Wronskian $W[\vec{y}_1(x), \ldots, \vec{y}_n(x)]$ of solutions $\vec{y}_1(x), \ldots, \vec{y}_n(x)$ is nonzero on (α, β) if and only if the solutions are linearly independent. It's *very important* here that the functions we are considering are solutions to the differential equation $\vec{y}(x) = A\vec{y}(x)$. To demonstrate this, consider the following functions

$$\vec{y}_1(x) = \begin{pmatrix} 1 \\ x \end{pmatrix}, \quad \vec{y}_2(x) = \begin{pmatrix} e^x \\ xe^x \end{pmatrix}$$

- (a) Show that $W[\vec{y}_1(x), \vec{y}_2(x)]$ is identically 0
- (b) Despite this, show that $\vec{y}_1(x)$ and $\vec{y}_2(x)$ are actually linearly independent

• • • • • • • • •