

# Math 309 Lecture 6 and 7

## Constant Coefficient Homogeneous Linear Systems of ODEs

W.R. Casper

Department of Mathematics  
University of Washington

March 27, 2017

Plan for today:

- Complex Eigenvalues
- Stability of the Origin

Next time:

- Fundamental Matrix

# Outline

# Review of Results from Last Time

- the origin is *always* a fixed point of  $\vec{y}'(t) = Ay(t)$
- eg.  $\vec{y}(t) = \vec{0}$  is a constant solution of the equation
- how other solutions behave is based on the eigenvalues of  $A$ :
  - (a) if both eigenvalues of  $A$  are real and positive, then origin is an exponentially unstable node
  - (b) if both eigenvalues of  $A$  are real and negative, then origin is an exponentially stable node
  - (c) if both eigenvalues of  $A$  are mixed sign, then origin is a saddle point
  - (d) what about when the eigenvalues of  $A$  are complex?

# Spirally Slope Fields

- slope fields for complex eigenvalues are characterized by spiral patterns
- for example:


$$A = \begin{pmatrix} -1/2 & 1 \\ -1 & -1/2 \end{pmatrix}$$

- characteristic polynomial is

$$p_A(x) = \det(A - xI) = x^2 + x + \frac{5}{4}$$

- eigenvalues of  $A$  are  $-(1/2) \pm i$


# Complex Eigenvalues: $-(1/2) \pm i$



# Behavior of the Origin


- suppose that  $A$  has complex eigenvalues
  - they come in conjugate pairs!  $\lambda_1 = a + ib$ ,  $\lambda_2 = a - ib$
  - the origin is *always* a fixed point of  $\vec{y}'(t) = Ay(t)$
  - whether our ship moves toward or away depends on value of  $a$
- (a) if  $a$  is positive, move away
- (b) if  $a$  is negative, move toward
- (c) if  $a$  is zero, circle around

# Complex Eigenvalues: $(1/2) \pm i$





# Complex Eigenvalues: $\pm i$



# What about General Solutions?

## Question

How do we find the general solution in the case that  $A$  has complex eigenvalues?

- use Euler's definition!

$$e^{i\theta} = \cos(\theta) + i \sin(\theta)$$

- we can then take our eigenvalue solutions and write them as linear combinations of *real* solutions

## Question

Find the general solution of the equation

$$\vec{y}'(x) = A\vec{y}, \quad A = \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}$$

- first we find the eigenvalues:  $-(1/2) \pm i$
- then we find the corresponding eigenspaces:

$$E_{-(1/2)+i} = \text{span} \left\{ \begin{pmatrix} 1 \\ i \end{pmatrix} \right\} \quad E_{-(1/2)-i} = \text{span} \left\{ \begin{pmatrix} 1 \\ -i \end{pmatrix} \right\}$$

# Example

- from this we get two (complex) solutions

$$\vec{y}_1(t) = \begin{pmatrix} 1 \\ i \end{pmatrix} e^{(-1/2+i)t} \quad \vec{y}_2(t) = \begin{pmatrix} 1 \\ -i \end{pmatrix} e^{(-1/2-i)t}$$

- by the superposition principle we get the family of solutions:

$$\begin{aligned} \vec{y}(t) &= c_1 \begin{pmatrix} 1 \\ i \end{pmatrix} e^{(-1/2+i)t} + c_2 \begin{pmatrix} 1 \\ -i \end{pmatrix} e^{(-1/2-i)t} \\ &= c_1 e^{-t/2} \begin{pmatrix} \cos(t) + i \sin(t) \\ i \cos(t) - \sin(t) \end{pmatrix} + c_2 e^{-t/2} \begin{pmatrix} \cos(t) - i \sin(t) \\ -i \cos(t) - \sin(t) \end{pmatrix} \\ &= e^{-t/2} \begin{pmatrix} (c_1 + c_2) \cos(t) + i(c_1 - c_2) \sin(t) \\ i(c_1 - c_2) \cos(t) - (c_1 + c_2) \sin(t) \end{pmatrix} \\ &= e^{-t/2} \begin{pmatrix} b_1 \cos(t) + b_2 \sin(t) \\ b_2 \cos(t) - b_1 \sin(t) \end{pmatrix} \\ &= b_1 e^{-t/2} \begin{pmatrix} \cos(t) \\ -\sin(t) \end{pmatrix} + b_2 e^{-t/2} \begin{pmatrix} \sin(t) \\ \cos(t) \end{pmatrix} \end{aligned}$$

# Correspondence

- the stability of the origin depends on the eigenvalues of the matrix  $A$
- five possibilities:
  - (1) all eigenvalues are real and negative  $\rightarrow$  origin is exponentially stable (sink)
  - (2) all of the eigenvalues are real and positive  $\rightarrow$  origin is unstable (source)
  - (3) eigenvalues are real and opposite-signed  $\rightarrow$  origin is saddle
  - (4) both of the eigenvalues are complex with positive real component  $\rightarrow$  origin is spirally unstable (spiral source)
  - (5) both of the eigenvalues are complex with nonpositive real component  $\rightarrow$  origin is spirally stable (spiral sink)

# Alternative Correspondence

- the stability may also be classified based on the determinant and trace of  $A$
- this is because

$$p_A(x) = x^2 - \operatorname{tr}(A)x + \det(A).$$

- five possibilities:
  - (1)  $\operatorname{tr}(A)^2 > 4 \det(A)$  and  $\det(A) < 0 \rightarrow$  saddle
  - (2)  $\operatorname{tr}(A)^2 > 4 \det(A)$ ,  $\det(A) > 0$ , and  $\operatorname{tr}(A) > 0 \rightarrow$  unstable
  - (3)  $\operatorname{tr}(A)^2 > 4 \det(A)$ ,  $\det(A) > 0$ , and  $\operatorname{tr}(A) < 0 \rightarrow$  exponentially stable
  - (4)  $\operatorname{tr}(A)^2 < 4 \det(A)$  and  $\operatorname{tr}(A) \leq 0 \rightarrow$  spirally stable
  - (5)  $\operatorname{tr}(A)^2 < 4 \det(A)$  and  $\operatorname{tr}(A) > 0 \rightarrow$  spirally unstable

# Stability Picture

*Principle - diagram: Classification of phase portraits in  $(\det A, \text{Tr} A)$ -plane*




Figure: Picture of Classification (DR Hundley, Whitman College)

# Summary!

What we did today:

- Complex Eigenvalues
- Stability of the Origin

Plan for next time:

- Fundamental Matrix